

CONTENTS

(Abstracts/contents lists published in: Chem. Abstr.; Curr. Contents: Phys., Chem. & Earth Sci.; Life Sci. Full texts are incorporated in CJELSEVIER, a file in the Chemical Journals Online database which is available on STN® International.)

Roles of potassium ions, acetyl and L-glyceryl groups in native gellan double helix: an X-ray study R. Chandrasekaran, A. Radha, and V. G. Thailambal (West Lafayette, IN, U.S.A.) . . .	1
E.s.r. study of the alkaline degradation of disaccharides in methyl sulfoxide P. Calle, A. Sanchez, and C. Sieiro (Madrid, Spain)	19
4-Thiocellooligosaccharides. Their synthesis and use as ligands for the separation of cellobiohydrolases of <i>Trichoderma reesei</i> by affinity chromatography C. Orgeret, E. Seillier, C. Gautier, J. Defaye, and H. Driguez (Grenoble, France)	29
Novel types of acyclic C-nucleoside analogues via the reaction of hydrogen bromide in acetic acid with L-threo-(glycerol-1-yl)pyrazolinediones H. A. Hamid, N. Rashed, Y. El Kilany, and E. S. H. El Ashry (Alexandria, Egypt)	41
Synthesis, structure, and transformations of 1,2,4-phosphites of α -D-xylopyranose E. E. Nifant'ev, M. P. Koroteev, V. A. Sychev, A. R. Bekker, L. K. Vasyanina, and N. N. Nevskii (Moscow, U.S.S.R.)	49
Spacer-modified disaccharide and pseudo-trisaccharide methyl glycosides that mimic maltotriose, as competitive inhibitors for pancreatic α -amylase: a demonstration of the "clustering effect" M. Blanc-Muesser, L. Vigne, H. Driguez (Grenoble, France), J. Lehmann, J. Steck, and K. Urbahns (Freiburg, Germany)	59
Preparation of higher-carbon sugars by stereoselective osmylation of related allylic alcohols S. Jarosz (Warsaw, Poland)	73
Synthesis of a tetrasaccharide acceptor for use in the assay of UDP-GlcNAc: β -D-Galp-(1 \rightarrow 4)- β -D-GlcNAc (GlcNAc to Gal) β (1 \rightarrow 3)-N-acetylglucosaminyltransferase activity and the pentasaccharide product that would be formed by its enzymic glycosylation G. Srivastava and O. Hindsgaul (Edmonton, Alta., Canada)	83
Synthesis and conformation of dihistamine derivatives of cyclomaltoheptaose (β -cyclodextrin) V. Cucinotta, F. D'Alessandro, G. Impellizzeri, and G. Vecchio (Catania, Italy)	95
Total stereo- and enantio-selective synthesis of 2,3-dideoxy-3-C-methylene-D-glycero-pentose and its ethyl furanosides Yu. E. Raifeld, G. Ya. Vid, I. E. Mikerin, B. M. Arshava, and A. A. Nikitenko (Moscow, U.S.S.R.)	103
Synthesis of nigero-oligosaccharides K. Takeo, S. Kitamura, and Y. Murata (Kyoto, Japan)	111
Synthesis of sialic acid-containing nucleotides sugars: CMP-sialic acid analogs K. Ikeda, Y. Nagao, and K. Achiwa (Shizuoka, Japan)	123

<i>In situ</i> synthesis of ferrites in lignocellulosics R. H. Marchessault, S. Ricard, and P. Rioux (Montreal, Que., Canada)	133
A synthetic study of methyl 3-deoxy-3-fluoro- α -D-glucopyranosides from methyl 2,3-anhydro- α -D-allopyranosides, and synthesis of 3'-deoxy-3'-fluorokanamycin A and 3'-chloro-3'-deoxykanamycin A E. Umemura, T. Tsuchiya, Y. Kobayashi, and K. Tanaka (Kawasaki, Japan)	141
Inclusion complexes of some non-ionic surfactants with cyclomalto-oligosaccharides T. Cserhati and J. Szejtli (Budapest, Hungary)	165
Synthesis of an S-(α -sialosyl)-(2 \rightarrow 9)-O-(α -sialosyl)-(2 \rightarrow 3')- β -lactosylceramide A. Hasegawa, H. Ogawa, H. Ishida, and M. Kiso (Gifu, Japan)	175
Synthesis of a position isomer of ganglioside GD ₃ having an α -Neu5Ac-(2 \rightarrow 9)- α -Neu5Ac linkage A. Hasegawa, H. Ogawa, and M. Kiso (Gifu, Japan)	185
Application of cyclic sulfates in the synthesis of 6-deoxy-D-manno-heptopyranose derivatives P. A. M. van der Klein and J. H. van Boom (Leiden, The Netherlands)	193
The distribution of carboxymethyl groups in O-(carboxymethyl)ated (1 \rightarrow 3)- β -D-glucans and (1 \rightarrow 3)- α -D-glucans S. Ukai, I. Yoshida, A. Honda, K. Nagai, and T. Kiho (Gifu, Japan)	201
Structure of a heteroglycan isolated from the fungus <i>Omphalia lapidescens</i> K. Saito, M. Nishijima, and T. Miyazaki (Tokyo, Japan)	209
New investigations of the structure of grape arabinogalactan-protein L. Saulnier, J.-M. Brillouet, M. Moutounet (Montpellier, France), C. Herve du Penhoat, and V. Michon (Paris, France)	219
Structure and antitumor activity of a branched (1 \rightarrow 3)- β -D-glucan from the alkaline extract of <i>Amanita muscaria</i> T. Kiho, M. Katsuragawa, K. Nagai, S. Ukai (Gifu, Japan), and M. Haga (Hokkaido, Japan)	237
Structure of the extracellular polysaccharide from slime-forming <i>Lactococcus lactis</i> subsp. <i>cremoris</i> SBT 0495 H. Nakajima, T. Hirota (Kawagoe, Japan), T. Toba, T. Itoh, and S. Adachi (Sendai, Japan)	245
The use of glycosides of 6- and 8-acylamino-4-methylumbelliferone in studies of the specificity and properties of human lysosomal glycolipid hydrolases G. Ya. Wiederschain, I. K. Kozlova, G. S. Ilyina, M. A. Mikhaylova, and E. M. Beyer (Moscow, U.S.S.R.)	255
<i>Notes</i>	
Crystal and molecular structure of 4,8-anhydro-3-deoxy-3-nitro-D-lyxo-D-glucos-nonitol P. Koll (Oldenburg, Germany), J. Kopf, M. Morf, B. Zimmer (Hamburg, Germany), M. Petrušova, and L. Petruš (Bratislava, Czechoslovakia)	273
Preparation and n.m.r. assignments of cellulose mixed esters regioselectively substituted by acetyl and propanoyl groups T. Iwata, J.-I. Azuma, K. Okamura (Kyoto, Japan), M. Muramoto (Osaka, Japan), and B. Chun (Kanagawa, Japan)	277
An n.m.r. study of the 1,3,4,6- and 1,2,3,6-tetra-acetates of β -D-allopyranose and the X-ray structure of the former E. Lee, P. Browne, P. McArdle, and D. Cunningham (Galway, Ireland)	285

Regioselective synthesis of α -L-fucosyl-containing disaccharides by use of α -L-fucosidases of various origins K. Ajisaka and M. Shirakabe (Odawara, Japan)	291
A new route to C-glycosyl compounds. Wittig-type reaction promoted by zinc L. Dheilly, C. Fréchou, D. Beaupère, R. Uzan, and G. Demailly (Amiens, France)	301
The use of chlorodimethylhexylsilane for protecting the hydroxyl groups in cyclomaltoheptaose (β -cyclodextrin) A. W. Coleman, P. Zhang, C.-C. Ling (Châtenay-Malabry, France), H. Parrot-Lopez, and H. Galons (Paris, France)	307
A facile synthesis of 4-O-allyl-D-xylopyranose and its use in the preparation of xylo-oligosaccharides K. Takeo, Y. Murata, and S. Kitamura (Kyoto, Japan)	311
A new synthesis of 2-deoxy- β -Kdo, a potent CMP-Kdo synthetase inhibitor H. Ohrui, M. Morita, and H. Meguro (Sendai, Japan)	319
Action of Cu(II) salts on osazones: side reactions of osotriazole formation H. El Khadem, P. Czubarow, and C. Sandstedt (Washington, DC, U.S.A.)	327
Structural studies of a polysaccharide from the seeds of <i>Nehumbo nucifera</i> S. Das, B. Ray, and P. K. Ghosal (Burdwan, India)	331
Location of the O-acetyl group in welan by the reductive-cleavage method J. D. Stankowski and S. G. Zeller (San Diego, CA, U.S.A.)	337
The effect of the content of D-mannuronic acid and L-guluronic acid blocks in alginates on the antitumor activity M. Fujihara and T. Nagumo (Kanagawa, Japan)	343
<i>Announcement</i>	c1
<i>Author index volume 224</i>	c3
<i>Subject index volume 224</i>	c5
<i>Contents volume 224</i>	c9